


# E05-009: MOLLER Status Update

Dustin McNulty

UMass

*mcnulty@jlab.org*


June 11, 2010



## E05-009: MOLLER Status Update


### Outline

- Intro: (fully approved at PAC34)
  - Moller Scattering, APV Measurement
  - Proposed Measurement Details
  - Goals and Motivation
- Experimental Setup/Design
  - Details: Beam, Target, Spectrometer
  - Simulation Studies
  - New Challenges
- Timeline and Status




## MOLLER Collaboration

- MOLLER – Measurement of Lepton-Lepton Electroweak Reaction
- ~100 authors from 30 institutions, with veterans from all the JLab parity violating experiments



## Moller Scattering, $A_{PV}$ Measurement




- Purely leptonic reaction provides clean probe of weak neutral current interactions via parity violating electroweak interference

$$A_{PV} = m_e E_{\text{lab}} \frac{G_F}{\sqrt{2}\pi\alpha} \frac{4\sin^2\theta}{(3 + \cos^2\theta)^2} Q_W^e, \quad (1)$$


$$Q_W^e \equiv 4 \cdot g_V^e \cdot g_A^e = (1 - 4\sin^2\theta_W) \quad (2)$$


- $e^-_{\text{beam}}$ : 11GeV, 85 $\mu$ A, 85% polarization  
 $\rightarrow \langle Q^2 \rangle = 0.0056 \text{ (GeV/c)}^2, \langle A_{PV} \rangle = 35.6 \text{ ppb}$
- For 38 week run:  $\delta(A_{PV}) = 0.74 \text{ ppb}, \delta(Q_W^e) = \pm 2.1(\text{stat}) \pm 1.0(\text{syst}):$   
 $\rightarrow \delta(\theta_W) = \pm 0.00026(\text{stat}) \pm 0.00012(\text{syst}) \sim 0.1\% \text{ precision!}$




## Physics Motivation: $\sin^2\theta_W$ , the Higgs Mass, and Beyond the Standard Model


- World data avg:  $\sin^2\theta_W = 0.23122(17)$   
 $\Rightarrow m_H = 89^{+38}_{-28} \text{ GeV}$   
(favors SUSY, rules out Technicolor)
- Avg dominated by two measurements separated by  $3\sigma$ :  
 $\rightarrow A_l(\text{SLD}) : 0.2310(3), \Rightarrow m_H = 35^{+26}_{-17} \text{ GeV}$   
rules out SM!  
 $\rightarrow A_{fb}^{0,1} : 0.2322(3), \Rightarrow m_H = 480^{+350}_{-230} \text{ GeV}$   
rules out SUSY, favors Technicolor
- Proposed measurement precise enough to effect the central value of  $\sin^2\theta_W$  and its implications for  $m_H$





## The Search for the Higgs





## Establishing Limits for New Contact Interactions (Off the Z Resonance)

Important component of indirect signatures for "new physics"



*consider*


$$\begin{aligned} & \text{Feynman diagram: } X \text{ (green wavy line)} \rightarrow \text{Contact interaction (cross with green dot)} \\ & \text{coupling: } A_X \propto \frac{1}{Q^2 - M_X^2} \\ & \sim \frac{4\pi}{\Lambda^2} \end{aligned}$$

$Q^2 \sim M_Z^2$     **on resonance:**  $A_Z$  *imaginary*     $\Rightarrow A_Z^2 \left[ 1 + \frac{A_X^2}{A_Z^2} \right]$

**no interference!**


- Proposed Measurement will reach  $\sim 7.5$  TeV interaction scale
- Best current limits on  $4e^-$  contact interac. come from LEP, LEPII:  
 $\Lambda/g \sim 5$  TeV, but insensitive to  $|g_{RR}^2 - g_{LL}^2|$
- Near the Z resonance, new physics interactions (e.g.  $Z'_X$  exchange)  
don't visibly mix with standard model  $A_Z$  (Collider Experiments)
- This underscores importance of low energy measurements of  $Q_W^e$ :  
E158, Qweak, PVDIS, and 12GeV Moller

Current and Future  $\sin^2 \theta_W$  Measurements




## Experimental Setup/Design



### Layout in Hall A




- Long and skinny design (~30m from target to detector)
- 150cm  $\text{IH}_2$  target
- Novel two toroid spectrometer design (prebender and hybrid) with full azimuthal acceptance
- Flux integrating detector ring with azimuthal and radial segmentation




## Optimized Spectrometer ( $\sim 100\%$ Acceptance)





All of those rays of  $\theta_{CM}=[90,120]$  that you don't get here...




... are collected as  $\theta_{CM}=[60,90]$  over here!




## Toroid Design Concept





--Two Toroid design facilitates signal and bkgd separation while focusing Moller events onto narrow detector ring







### Optics Raytrace




--Defocusing effects results in population of full azimuth




## Main Detector Reference Design



- Rad-hard flux, integrating detectors
- Radial segmentation for systematic checks (backgrounds)
- Azimuthal segmentation for systematic checks (e.g. parity conserving  $\cos(\phi)$  asym, azimuthal defocusing, beam sensitivities, backgrounds, etc.)
- Ancillary detectors (not shown): Tracking, pion, and lumi




Side View



## New Challenges

- 150GHz total detected Moller event rate
  - Must flip pockels cell at  $\sim 2\text{kHz}$
  - 80ppm pulse-to-pulse statistical fluctuations
    - Electronic noise and density fluctuations  $< 10^{-5}$
    - Pulse-to-pulse beam monitoring res. a few microns at 1kHz
- 0.5nm/0.05nrad control of beam on target
  - Requires improvement on control of pol. src. laser transport
  - Improved methods of “slow helicity reversal” (double wien)
- Target requires  $\sim 5\text{kW}$  of cooling power at  $85\mu\text{A}$   $I_{\text{beam}}$
- Full azimuthal acceptance with  $\theta_{\text{lab}}$  between 5 and 17mrad
  - Aggressive spectrometer design
  - Complex collimation and shielding issues
- Robust and redundant 0.4% beam polarimetry
  - Plan to pursue both Compton and atomic Hydrogen techniques




## Timeline and Status


- JLab PAC 34 - full approval - strong endorsement

“The proposed physics reach is outstanding and capable of making this effort a flagship experiment at JLab. The PAC believes the mission of this experiment... is so important that the Laboratory should make every effort to support the securing of the resources required”

- Detailed cost estimates and R&D plan formulation underway
- Working with lab management to prepare funding request (DOE, NSF, and international funding agencies)
- First review (JLab director's review) took place last January – Addressed charges: Physics case, and experimental approach
- Construction/Installation: 2012 - 2015
- Experiment likely to have two running periods: The first 6 months, and second 2 years





## Extra Slides





## Complimentary Measurement to LHC

$Z'$  Leptonic Couplings,  $M_{Z'} = 1.5$  TeV






## Transverse Asymmetry Measurement




Average transverse asymmetry




-- $A_T \sim 12$  parts per million ... 3 orders of magnitude bigger than  $A_{PV}$


--We must be sure that this averages to a negligible contribution!




## Magnet Design

# TOSCA







## Main Detector Reference Design



# MOLLER Collaboration

Jefferson Lab Hall A





## Spectrometer Beamline Draft

